

Aberystwyth University

Geographically extensive larval surveys reveal an unexpected scarcity of primary vector mosquitoes in a region of persistent malaria transmission in western Zambia

Cross, Donall; Thomas, Chris; McKeown, Niall; Siazuyu, Vincent; Healey, Amy; Willis, Tom; Singini, Douglas; Liywalii, Francis; Silumesii, Andrew; Sakala, Jacob; Smith, Mark ; Macklin, Mark; Hardy, Andy; Shaw, Paul

Published in:
Parasites & Vectors

DOI:
[10.1186/s13071-020-04540-1](https://doi.org/10.1186/s13071-020-04540-1)

Publication date:
2021

Citation for published version (APA):
Cross, D., Thomas, C., McKeown, N., Siazuyu, V., Healey, A., Willis, T., Singini, D., Liywalii, F., Silumesii, A., Sakala, J., Smith, M., Macklin, M., Hardy, A., & Shaw, P. (2021). Geographically extensive larval surveys reveal an unexpected scarcity of primary vector mosquitoes in a region of persistent malaria transmission in western Zambia. *Parasites & Vectors*, 14(1), [91]. <https://doi.org/10.1186/s13071-020-04540-1>

Document License CC BY

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Cross *et al.* Geographically extensive larval surveys reveal an unexpected scarcity of primary vector mosquitoes in a region of persistent malaria transmission in western Zambia




Figure S1: Maximum Likelihood phylogenetic tree of cytochrome c oxidase I (COI) sequences from *Anopheles* larvae sampled in western Zambia. Consensus tree constructed in MEGA X from a 318bp fragment of the COI region of mitochondrial DNA, using 100 bootstrap replicates. Sequences from this paper have been labelled with taxon assignment used in this paper. Published reference sequences are labelled with NCBI nt accession number and species ID; suffix denotes source paper. ‘--L’ denotes Lobo *et al.* (2015); ‘--SL’ denotes St Laurent *et al.* (2016).

St Laurent B, Cooke M, Krishnankutty SM, Asih P, Mueller JD, Kahindi S, et al. Molecular characterization reveals diverse and unknown malaria vectors in the western Kenyan highlands. *Am J Trop Med Hyg.* 2016;94:327-35; doi: 10.4269/ajtmh.15-0562.

Lobo NF, St Laurent B, Sikaala CH, Hamainza B, Chanda J, Chinula D, et al. Unexpected diversity of *Anopheles* species in eastern Zambia: implications for evaluating vector behavior and interventions using molecular tools. *Sci Rep-Uk.* 2015;5; doi: 10.1038/srep17952.